

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

0 310 949  
A2

(12)

# EUROPEAN PATENT APPLICATION

(21) Application number: 88116110.3

(51) Int. Cl.4: C12P 13/06 , C12P 41/00 ,  
/(C12P13/06,C12R1:13)

(22) Date of filing: 29.09.88

(30) Priority: 07.10.87 JP 253127/87

(43) Date of publication of application:  
12.04.89 Bulletin 89/15

(94) Designated Contracting States:  
CH DE FR GB IT LI SE

(71) Applicant: TORAY INDUSTRIES, INC.  
2, Nihonbashi-Muromachi 2-chome Chuo-ku  
Tokyo 103(JP)

(72) Inventor: Takeuchi, Masae  
16, Tenjin Kitano-cho  
Konan-shi Aichi 484(JP)  
Inventor: Yonehara, Tetsu  
7-5, Sanjou 2-chome Minami-ku  
Nagoya-shi Aichi, 457(JP)

(74) Representative: Kador & Partner  
Corneliusstrasse 15  
D-8000 München 5(DE)

(54) Process for producing D-alanine.

(57) It is possible to produce and accumulate D-alanine selectively and to improve the amount of production and accumulation of D-alanine by cultivating a microorganism having both an ability to produce D-alanine and a resistance to D-cycloserine and belonging to the genus Brevibacterium.




Figure 1

Xerox Copy Centre

EP 0 310 949 A2

## PROCESS FOR PRODUCING D-ALANINE

Background of the Invention(1) Field of the Invention

The present invention relates to an industrially favorable process for producing D-alanine.

D-alanine is an unnatural amino acid and it is a useful compound for a reagent as it is or for a raw material for synthesizing peptides and so on. Accordingly, the demand has been recently increasing.

(2) Description of the Prior Art

A number of processes for producing DL-alanine by means of a fermentation process using a microorganism belonging to the genus Brevibacterium have been hitherto known. An improved process for producing DL-alanine by using a microorganism belonging to the genus Brevibacterium and having sensitivity against threonine and methionine, also has been known (J.Gen. Appl. Microbiol. 1971, 17(2), 169-72).

However, it is inevitable to use a certain optical resolution operation for obtaining only D-alanine, because L-alanine is also produced by these process.

Namely, it has never been achieved yet to produce D-alanine by fermentation with low cost, simply and industrially favorably.

Summary of the Invention

One of the purposes of the present invention is to offer a process for producing D-alanine by means of an optically selective fermentation method.

The second purpose of the present invention is to offer a process for increasing the accumulative concentration of D-alanine by fermentation using one microorganism.

The third purpose of the present invention is to offer a process for producing D-alanine without using an optical resolution operation.

The fourth purpose of the present invention is to offer a process for producing D-alanine with low cost and simply.

The final purpose of the present invention is to offer an industrially favorable process for producing D-alanine.

Other and further objects, features and advantages of the present invention will appear more fully in the following description.

These purposes can be achieved by a process for producing D-alanine characterized by cultivating a microorganism, which belongs to the genus Brevibacterium and has an ability to produce D-alanine and a resistance to D-cycloserine, producing and accumulating thereby D-alanine in a culture medium, and obtaining D-alanine from the culture broth.

Brief Description of the Drawings

Figure 1 shows the quantities of D- and L- alanine, respectively, accumulated in the culture medium with the passage of cultivation time.

Figure 2 shows the quantities of D- and L- alanine, respectively, accumulated in the culture medium with the elapse of cultivation time.

Description of the Preferred Embodiments

The microorganisms used in the present invention are those microorganisms that have an ability to produce D-alanine and a resistance to D-cycloserine and belong to the genus Brevibacterium. Whenever any microorganism belonging to the genus Brevibacterium has these characteristics, it can be included in the range of the present invention even if it has other auxotrophy and drug resistance.

Microorganisms which belong to the genus Brevibacterium and have resistance to D-cycloserine can produce and accumulate D-alanine selectively with high accumulation quantity.

As the representative samples of the microorganisms used in the present invention, for example, Brevibacterium lactofermentum DCSR-26, Brevibacterium lactofermentum DCSR17-2 and Brevibacterium flavum DCSR 2-2 can be cited. Brevibacterium lactofermentum DCSR-26 is derived from Brevibacterium lactofermentum ATCC 13869 (requiring biotin for the growth) and it is a variant having resistance to D-cycloserine. Brevibacterium lactofermentum DCSR17-2 is derived from Brevibacterium lactofermentum DCSR-26 and it is a variant having a further higher resistance to D-cycloserine. Brevibacterium flavum DCSR 2-2 is derived from Brevibacterium flavum ATCC 13826 and it is a variant having a resistance to D-cycloserine.

Brevibacterium lactofermentum DCSR-26 was deposited to Fermentation Research Institute, Agency of Industrial Science and Technology in Japan (abbreviated as FRI thereafter) on September 24, 1987 and a registration No. FERM BP-2023 was conferred on it. Brevibacterium lactofermentum DCSR 17-2 was deposited to FRI in Japan on March 9, 1988 and a registration No. FERM BP-2024 was conferred on it. Brevibacterium flavum DCSR 2-2 was deposited to FRI in Japan on September 12, 1988 and a registration No. FERM BP-2047 was conferred on it.

FERM BP number is the registration number of the deposition to FRI based on Budapest Treaty. All restriction upon availability to the public will be irrevocably removed upon granting of the patent and the deposit will be maintained in a public depository for a period of 30 years or 5 years after the last request or for the effective life of the patent, which is longer.

FRI is located in 1-1, Higashi 1-Chome, Tsukuba, Ibaraki ken, Japan.

Derivation of the variant can be relatively easily carried out by means of an ordinary mutative treatment. Namely, to obtain variants having resistance to D-cycloserine, parent strains are either irradiated by ultraviolet or treated with mutagenic agents (for example, N-methyl-N'-nitro-N nitrosoguanidine, ethyl methanesulfonate and so on) and strains which are capable of growing distinguishably in comparison with the parent strain can be selected from the culture plates contained D-cycloserine in such a concentration that the parent stain can not grow sufficiently.

D-cycloserine resistant strain in the present invention means the strain gaining stronger resistance than that the parent strain has. The relative growth degree of the D-cycloserine resistant strain is preferably 50% or more when the cultivation is carried out in a culture medium containing D-cycloserine with such a concentration that the relative growth degree of the parent strain becomes 40% or less. Here, the relative growth degree is shown by the percentage of the optical density measured at 660nm of the culture broth containing no D-cycloserine.

As the culture medium used in the present invention, various nutrient resources being widely used in cultivation of microorganisms can be used. For example, as the carbon source, sugars such as glucose, molasses, hydrolysate of starch and so on, organic acids such as acetic acid, benzoic acid and so on, alcohols such as ethanol and so on can be used and as the nitrogen source, ammonium sulfate, ammonium nitrate, ammonium chloride, ammonium phosphate, urea, ammonia and so on can be used. Depending upon the kind of inorganic ammonium salts, inorganic salts such as phosphate, calcium carbonate and so on are required to use with them. Moreover, to improve the growth of microorganisms and to accumulate large amount of D-alanine, it is preferable to add, for example, organic nitrogen sources, vitamins, a small amount of metal ions and so on in the above described culture medium. Consequently, the purpose can be sufficiently achieved by adding usually cheap natural products such as soybean hydrolysate and corn steep liquor.

In the present invention, it is preferable to add D,L-alanine containing L-alanine with any ratio in the culture medium. As the additive D,L-alanine containing L-alanine with any ratio, not only a racemic modification of D,L-alanine but also pure L-alanine can be used. The time of the addition is not specifically restricted during the cultivation, but the early or intermediate phase of the cultivation is the most effective. By adding D,L-alanine, the ability of D-alanine accumulation by the microorganism used in the present invention can be improved and L-alanine is converted into D-alanine by the action of the microorganism. As the result, the amount of production and accumulation of D-alanine is surprisingly improved. The amount of

D,L-alanine to be added is not specifically restricted. The larger the amount added, the higher the amount of accumulation of D-alanine. Usually, it is 100g/l or less.

In the present invention, the above described cultivation of the microorganism is preferably carried out in an aerobic condition such as shaking or stirring with aeration. The cultivation temperature is usually in the range of 20-40 °C and especially around 30 °C is preferable. The pH of the medium is usually at 5-9 and it is preferable to keep it at about neutral for obtaining high yield. Thus, after cultivation for several days, a remarkable amount of D-alanine is produced and accumulated in the culture broth. After the cultivation is finished, D-alanine produced can be easily obtained by well known isolation and purification procedures such as ion exchange, adsorption and precipitation procedures without using an optical resolution method.

The present invention will be more clearly understood with reference to the following Experiments and Examples.

#### Experiment 1

##### A. (Separation of D-cycloserine resistant strains)

Cells of *Brevibacterium lactofermentum* ATCC 13869 (requiring biotin for the growth) were treated with N-methyl-N'-nitro-N-nitrosoguanidine (300 µg/ml, at 30 °C for 10 minutes) by means of an usual procedure and then spread on the complete synthetic agar plate containing 50 mg/l of D-cycloserine (the complete synthetic agar plate : 2% of glucose, 2% of agar, 1% of ammonium sulfate, 0.1% of potassium hydrogen-phosphate, 0.04% of 7 hydrate of magnesium sulfate, 0.05% of sodium chloride, 0.25% of urea, 10 mg/l of 7 hydrate of ferrous sulfate, 10 mg/l of 4 hydrate of manganese (II) sulfate and 50 µg/l of biotin). Then, the incubation was carried out at 30 °C for 5-7 days and large colonies appeared were separated by means of colony picking to select a D-cycloserine resistant strain (*Brevibacterium lactofermentum* DCSR-26).

Another D-cycloserine resistant strain (*Brevibacterium flavum* DCSR 2-2) was obtained in the same way as the above described procedures such as mutation, cultivation and separation of the strain except that *Brevibacterium lactofermentum* ATCC 13869 was replaced with *Brevibacterium flavum* ATCC 13826.

Moreover, cells of *Brevibacterium lactofermentum* DCSR-26 were treated with N-methyl-N'-nitro-N-nitrosoguanidine (300 µg/ml, at 30 °C for 10 minutes) by means of an usual procedure and then spread on the complete synthetic agar plate containing 100 mg/l of D-cycloserine (the complete synthetic medium : 2% of glucose, 2% of agar, 1% of ammonium sulfate, 0.1% of potassium hydrogen-phosphate, 0.04% of 7 hydrate of magnesium sulfate, 0.05% of sodium chloride, 0.25% of urea, 10 mg/l of 7 hydrate of ferrous sulfate, 10 mg/l of 4 hydrate of manganese (II) sulfate and 50 µg/l of biotin). Then, the cultivation was carried out at 30 °C for 5-7 days and large colonies appeared were separated by means of colony picking to select a D-cycloserine highly resistant strain (*Brevibacterium lactofermentum* DCSR 17-2).

##### B. (The degree of resistance of D-cycloserine resistant strains)

Each strain shown in Table 1 was cultivated with shaking by using a bouillon liquid medium at 30 °C for 16 hours and the cells grown were harvested and washed well with sterilized physiological saline. This cells suspension was inoculated in 5 ml of minimal media contained D-cycloserine in concentrations of 0, 10 and 50 mg/l, respectively, (the minimal medium containing 2% of glucose, 1% of ammonium sulfate, 0.1% of potassium hydrogen-phosphate, 0.04% of 7 hydrate of magnesium sulfate, 0.05% of sodium chloride, 0.25% of urea, 10 mg/l of 7 hydrate of ferrous sulfate, 10 mg/l of 4 hydrate of manganese (II) sulfate and 50 µg/l of biotin) and cultivations were continued for 64 hours at 30 °C to investigate the degree of growth of each strain. The results were shown in Table 1.

It was clear from the data shown in Table 1 that D-cycloserine resistant strains used in the present invention (*Brevibacterium lactofermentum* DCSR-26, *Brevibacterium lactofermentum* 17-2 and *Brevibacterium flavum* DCSR 2-2) grew without obstruction by D-cycloserine and were resistant to D-cycloserine in comparison with the respective parent strains, *Brevibacterium lactofermentum* ATCC 13869 and *Brevibacterium flavum* ATCC 13826.

Table 1

| Strain                          | Relative growth degree (%)                  |         |         |
|---------------------------------|---|---------|---------|
|                                 | Concentration of D-cycloserine added (mg/l) |         |         |
|                                 | 0   | 10      | 50      |
| Parent strain                   |   |         |         |
| <u>Brevibacterium</u>           | 100   | 36.7    | 0       |
| <u>lactofermentum</u>           | (60.0)                                      | (22.0)  | (0)     |
| ATCC 13869                      |   |         |         |
| Strain of the present invention |   |         |         |
| <u>Brevibacterium</u>           | 100   | 78.4    | 61.8    |
| <u>lactofermentum</u>           | (87.0)                                      | (68.2)  | (53.8)  |
| DSCR-26                         |   |         |         |
| Strain of the present invention |   |         |         |
| <u>Brevibacterium</u>           | 100   | —       | 109     |
| <u>lactofermentum</u>           | (92.9)                                      |         | (101.7) |
| DCSR 17-2                       |   |         |         |
| Parent strain                   |   |         |         |
| <u>Brevibacterium</u>           | 100   | 87      | 82      |
| <u>flavum</u>                   | (125.2)                                     | (109.0) | (103.0) |
| ATCC 13826                      |   |         |         |
| Strain of the present invention |   |         |         |
| <u>Brevibacterium</u>           | 100   | 103     | 107     |
| <u>flavum</u> DCSR 2-2          | (103.1)                                     | (106.6) | (110.8) |

Notice: Optical densities at 660 nm of culture broths were shown in ( ). The relative growth degree was calculated by using the optical density of the culture broth where no D-cycloserine was added as 100%.

Example 1

15

20

25

Each 40 ml of fermentation medium (pH 7.25) containing 10% of glucose, 1.8% ammonium sulfate, 0.05% of di-potassium hydrogenphosphate, 0.025% of 7 hydrate of magnesium sulfate, 0.001% of zinc chloride, 2% of soybean hydrolysate (Ajinomoto, Tokyo, Japan), 3% of calcium carbonate and 30  $\mu$ g/l of biotin was poured into 1-l Erlenmeyer flask and sterilized in an autoclave at 120°C for 10 minutes to obtain fermentation medium. Brevibacterium lactofermentum DCSR-26 and its parent strain, Brevibacterium lactofermentum ATCC 13869 were inoculated in a bouillon liquid medium respectively and cultivated with shaking at 30°C for 5 days. 1% of the above described each culture medium was put into the above described fermentation medium respectively. After shaking at 30°C for 5 days, D-alanine was produced in each culture medium as shown in Table 2.

Table 2

|   |                       | Amount of<br>D-alanine<br>accumulated | Purity* |
|---|-----------------------|---------------------------------------|---------|
| Strain                                    |                       |                                       |         |
| Comparative<br>example                    | <u>Brevibacterium</u> |                                       |         |
|   | <u>lactofermentum</u> | 9.2 g/l                               | 76%     |
|   | ATCC 13869            |                                       |         |
| Example<br>of the<br>present<br>invention | <u>Brevibacterium</u> |                                       |         |
|   | <u>lactofermentum</u> | 13.2 g/l                              | 100%    |
|   | DCSR-26               |                                       |         |

$$\star \text{ Purity} = \frac{\text{D-alanine (g/l)}}{\text{Total alanine (g/l)}} \times 100\%$$

55 Total amount of alanine in the fermentation medium was analysed by means of an automatic amino acid analyzer. Analysis of D-alanine was carried out both by means of an enzymatic method using D-amino acid oxidase and by means of high performance liquid chromatography (HPLC) using a column for optical resolution (Sumitomo Chemical Industries, Co., Ltd OA-1000).

The Cells were removed by means of centrifugal separation from 1l of the fermentation broth of

Brevibacterium lactofermentum DCSR-26 to obtain supernatant, which was then decolorized with active carbon powder. The decolorized liquid thus obtained was passed through a column where a strong cation exchange resin "DIAION SK-1B" (H<sup>+</sup> type) was packed to adsorb D-alanine, which was eluted with 2N aqueous ammonia after washing with water. The eluted fractions of D-alanine were concentrated and ethanol was added to the concentrated liquid to obtain precipitated crystals. 9.5g of D-alanine crystals were obtained by recrystallizing with ethanol.

Specific rotation  $[\alpha]_D^{25} = -14.2^\circ$  (C=6, 1N-HCl)

#### 10 Example 2.

1 l of the fermentation medium described as in Example 1 without calcium carbonate were put into a jar fermenter and sterilized. Brevibacterium lactofermentum DCSR 17-2 was cultivated in the bouillon liquid medium at 30°C for 16 hours. Then, 40 ml of this culture medium was inoculated in the above described 1-l fermentation medium and fermentation was continued aerobically (1v.v.m., impeller speed : 600r.p.m.) for 3 days at 30°C to produce 14.8g/l of D-alanine in the broth. The relations between the cultivation time and the accumulated amount of alanine for 3 days were shown in Figure 1.

D-alanine was separated and obtained from 1 l of the culture broth by the same procedures as described in Example 1 and 11.8g of D-alanine crystals were thereby obtained.

20 Specific rotation  $[\alpha]_D^{25} = -14.2^\circ$  (C=6, 1N-HCl)

#### Example 3

25 40 ml of fermentation medium (pH 7.25) containing 10% of glucose, 3% of ammonium sulfate, 0.05% of potassium hydrogen-phosphate, 0.05% of dipotassium hydrogen-phosphate, 0.025% of 7 hydrate of magnesium sulfate, 2% of soybean hydrolysate, 3% of calcium carbonate and 30μg/l of biotin was poured into 1-l Erlenmeyer flask and sterilized in an autoclave at 120°C for 20 minutes to obtain culture medium. Brevibacterium lactofermentum DCSR 17-2 was inoculated in the fermentation medium thus obtained and cultivated at 30°C for 3 days. 20g/l of racemic D,L-alanine was therein added and the cultivation was furthermore carried out for 64 hours to produce and accumulate 48.8g/l of D-alanine in the culture broth as shown in Figure 2.

The supernatant liquid obtained by removing cells from 200 ml of this culture broth by means of centrifugal separation was decolorized with active carbon powder. The decolorized liquid was passed 35 through a column where a strong cation exchange resin "DIAION SK-1B" (H<sup>+</sup> type) was packed to adsorb D-alanine, which was eluted with 2N aqueous ammonia after washing with water. The eluted fractions of D-alanine were concentrated and ethanol was added to concentrated liquid to obtain precipitated crystals. 7.2g of D-alanine crystals were obtained by recrystallizing with ethanol.

Optical purity 99.4%

40 Specific rotation  $[\alpha]_D^{25} = -14.2^\circ$  (C=6, 1N-HCl)

Analysis of D-and L-types of alanine was carried out in the same was as described in Exampels 1.

#### Example 4

45 Each 40 ml of fermentation medium (pH 6.5) containing 10% of glucose, 2% of ammonium sulfate, 0.05% of potassium hydrogen-phosphate, 60μg/l of biotin, 1% of sodium chloride, 10 mg/l of 7 hydrate of ferrous sulfate, 20 mg/l of 4 hydrate of manganese (II) sulfate, 0.025% of 7 hydrate of magnesium sulfate and 2% of hot water extract of bean cake was poured into 1 l Erlenmeyer flask and sterilized in an autoclave at 120°C for 20 minutes to obtain culture medium. Brevibacterium flavum ATCC 13826 and 50 Brevibacterium flavum DCSR 2-2 were respectively inoculated in the fermentation media thus obtained and shaken at 30°C for 3 day. D-alanine was thereby produced in each culture broth as shown in Table 3.

Table 3

|   | Strain                                 | Amount of<br>D-alanine<br>accumulated | Purity | D/L<br>ratio* |
|---|--|---------------------------------------|--------|---------------|
| Comparative<br>example                    | <u>Brevibacterium</u><br><u>flavum</u> | 1.7 g/l                               | 85%    | 6.96          |
|   | ATCC 13826                             |                                       |        |               |
| Example<br>of the<br>present<br>invention | <u>Brevibacterium</u><br><u>flavum</u> | 1.9 g/l                               | 95%    | 13.4          |
|   | DCSR 2-2                               |                                       |        |               |

$$* \text{ D/L ratio} = \frac{\text{Amount of D-alanine}}{\text{Amount of L-alanine}}$$

As the D/L ratio was larger than 10 in the Example of the present invention, it was possible to separate D-alanine with almost no loss by means of recrystallization.

30

#### Claims

1. A process for producing D-alanine characterized by cultivating a microorganism which belongs to the  
35 genus Brevibacterium and has an ability to produce D-alanine and a resistance to D-cycloserine, producing and accumulating thereby D-alanine in a culture broth and obtaining D-alanine from the culture broth.
2. A process according to Claim 1, D,L-alanine containing L-alanine with any ratio is added in the culture medium.
3. A process according to Claim 1 or 2, the microorganism is a microorganism belonging to the genus  
40 Brevibacterium, the species lactofermentum or flavum.
4. A process according to Claim 3, the microorganisms are those derived from the strain ATCC 13869 and belonging to the genus Brevibacterium, the species lactofermentum.
5. A process according to Claim 3, the microorganisms are those derived from the strain ATCC 13826 and belonging to the genus Brevibacterium, the species flavum.

45

50

55






Figure 1




Figure 2

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 310 949**  
**A3**

(12)

# EUROPEAN PATENT APPLICATION

(21) Application number: 88116110.3

(51) Int. Cl. 4: **C12P 13/06** , **C12P 41/00** ,  
//(**C12P13/06,C12R1:13**)

(22) Date of filing: 29.09.88

(30) Priority: 07.10.87 JP 253127/87

(43) Date of publication of application:  
12.04.89 Bulletin 89/15

(84) Designated Contracting States:  
**CH DE FR GB IT LI SE**

(88) Date of deferred publication of the search report:  
02.11.89 Bulletin 89/44

(71) Applicant: **TORAY INDUSTRIES, INC.**  
**2, Nihonbashi-Muromachi 2-chome Chuo-ku**  
**Tokyo 103(JP)**

(72) Inventor: **Takeuchi, Masae**  
**16, Tenjin Kitano-cho**  
**Konan-shi Aichi 484(JP)**  
Inventor: **Yonehara, Tetsu**  
**7-5, Sanjou 2-chome Minami-ku**  
**Nagoya-shi Aichi, 457(JP)**

(74) Representative: **Kador & Partner**  
**Corneliusstrasse 15**  
**D-8000 München 5(DE)**

(54) **Process for producing D-alanine.**

(57) It is possible to produce and accumulate D-alanine selectively and to improve the amount of production and accumulation of D-alanine by cultivating a microorganism having both an ability to produce D-alanine and a resistance to D-cycloserine and belonging to the genus Brevibacterium.




Figure 1

Xerox Copy Centre

EP 0 310 949 A3



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number

EP 88 11 6110

| DOCUMENTS CONSIDERED TO BE RELEVANT  |  |  |  |
|--|--|--|--|
| Category   | Citation of document with indication, where appropriate, of relevant passages  | Relevant to claim                              | CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)         |
| Y  | CHEMICAL ABSTRACTS, vol. 67, no. 15, 9th October 1967, page 6700, abstract no. 71293s, Columbus, Ohio, US; R.H. REITZ et al.: "The biochemical mechanisms of resistance by streptococci to the antibiotics D-cycloserine and O-carbamyl-D-serine", & BIOCHEMISTRY 6(8), 2561-70(1967)<br>* Abstract *<br>---   | 1-5  | C 12 P 13/06 //<br>(C 12 P 13/06<br>C 12 R 1:13 )      |
| Y  | PATENT ABSTRACTS OF JAPAN, vol. 8, no. 232 (C-248)[1669], 25th October 1984; & JP-A-59 113 893 (NIPPON CARBIDE KOGYO K.K.) 30-06-1984<br>* Abstract *<br>---   | 1-5  |  |
| Y  | CHEMICAL ABSTRACTS, vol. 87, no. 7, 15th August 1977, page 200, abstract no. 49947z, Columbus, Ohio, US; V.L. CLARK et al.: "D-Cycloserine-induced alterations in the transport of D-alanine and glycine in Bacillus subtilis 168", & ANTIMICROB. AGENTS CHEMOTHER. 1977, 11(5), 877-80<br>* Abstract *<br>--- | 1-5  | TECHNICAL FIELDS<br>SEARCHED (Int. Cl.4)<br><br>C 12 P |
| A  | US-A-3 871 959 (ICHIRO CHIBATA)<br>---   |  |  |
| A  | BIOTECHNOLOGY OF AMINO ACID PRODUCTION, PROGRESS IN INDUSTRIAL MICROBIOLOGY, vol. 24, 1986, pages 3-13, Elsevier, Amsterdam, NL<br>---<br>-/-  |  |  |
| The present search report has been drawn up for all claims   |  |  |  |
| Place of search<br>THE HAGUE   |  | Date of completion of the search<br>04-08-1989 | Examiner<br>PULAZZINI A.F.R.                           |
| <b>CATEGORY OF CITED DOCUMENTS</b><br>X : particularly relevant if taken alone<br>Y : particularly relevant if combined with another document of the same category<br>A : technological background<br>O : non-written disclosure<br>P : intermediate document<br><br>T : theory or principle underlying the invention<br>E : earlier patent document, but published on, or after the filing date<br>D : document cited in the application<br>L : document cited for other reasons<br>.....<br>& : member of the same patent family, corresponding document |  |  |  |

EPO FORM 1503 03.82 (P0401)



EP 88 11 6110

| DOCUMENTS CONSIDERED TO BE RELEVANT   |   |   |  |
|---|---|---|--|
| Category  | Citation of document with indication, where appropriate, of relevant passages   | Relevant to claim   | CLASSIFICATION OF THE APPLICATION (Int. Cl. 4) |
| D, A  | J. GEN. APPL. MICROBIOL., vol. 17, no. 2, 1971, pages 169-172; K. SANO et al.: "Microbial production of alanine, L-proline, and L-valine by threonine- or methionine-sensitive mutants of <i>brevibacterium flavum</i> "<br>----- |   |  |
|   |   |   | TECHNICAL FIELDS SEARCHED (Int. Cl. 4)         |
|   |   |   |  |
| The present search report has been drawn up for all claims  |   |   |  |
| Place of search<br>THE HAGUE  |   | Date of completion of the search<br>04-08-1989  | Examiner<br>PULAZZINI A.F.R.                   |
| <b>CATEGORY OF CITED DOCUMENTS</b>  |   |   |  |
| X : particularly relevant if taken alone<br>Y : particularly relevant if combined with another document of the same category<br>A : technological background<br>O : non-written disclosure<br>P : intermediate document |   | T : theory or principle underlying the invention<br>E : earlier patent document, but published on, or after the filing date<br>D : document cited in the application<br>L : document cited for other reasons<br>-----<br>& : member of the same patent family, corresponding document |  |



11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000